

The process $e^+e^- \rightarrow \bar{t}c$ in topcolor-assisted technicolor models

Chongxing Yue^(a,b), Yuanben Dai^(b,c), Qingjun Xu^b, Guoli Liu^b

a: CCAST (World Laboratory) P.O. BOX 8730. B.J. 100080 P.R. China

b: College of Physics and Information Engineering,

Henan Normal University, Xinxiang 453002. P.R.China ^{*†}

c: Institute of Theoretical Physics, Academia Sinica, P.O. BOX 2735, B.J. 100080 P.R. China

February 1, 2008

Abstract

We consider the flavor changing neutral current(FCNC) process $e^+e^- \rightarrow \bar{t}c$ (or $t\bar{c}$) in the context of topcolor-assisted technicolor(TC2) models. We find that the contributions of charged scalars (technipions and top-pions) and topcolor gauge bosons are negligibly small. The neutral scalars (top-pion π_t^0 and top-Higgs h_t^0) can give significant contributions to this process. With reasonable values of the parameters in TC2 models, there will be several tens of the observable $\bar{t}c$ events to be generated in the future e^+e^- linear colliders.

PACS number(s): 14.65.Ha; 14.80.Cp

^{*}This work is supported by the National Natural Science Foundation of China(I9905004), the Excellent Youth Foundation of Henan Scientific Committee(9911), and Foundation of Henan Educational Committee.

[†]E-mail:cxyue@public.xxptt.ha.cn

It is widely believed that the top quark, with a mass of the order of the electroweak scale, will be a sensitive probe into physics beyond the standard model(SM).The properties of the top quark could reveal information about flavor physics, electroweak symmetry breaking(EWSB), as well as new physics beyond the SM[1]. Much attention has been paid to the studies of its production and decays. One of the most important tasks of high-energy linear e^+e^- colliders(LC) and hadron colliders is the careful investigation of the production of top quark. As the environment of the process $e^+e^- \rightarrow \bar{t}c$ (or $t\bar{c}$) is very clean, though the top quark has been found at Tevatron, the LC still potentially offer a complement to those of the hadron colliders studying the production of the top quark.

The search for flavor changing neutral current(FCNC) is one of the most interesting possibilities to test the SM, with the potential for either discovering or putting stringent bounds on new physics. In the SM, there are no FCNC at tree-level and at one-loop level they are GIM suppressed, which can offer a good place to test quantum effects of the fundamental quantum field theory. In models beyond SM, new particles may appear in the loop and have significant contributions to FCNC processes. Therefore, FCNC interactions give an ideal place to search for new physics. Any positive observation of FCNC couplings deviated from that in SM would unambiguously signal the presence of new physics. Thus, searching for FCNC is also one of the major goals of the next generation of high energy colliders, such as LC[2].

The FCNC vertices $tcV(V = \gamma, Z)$ can be probed either directly in the top-quark decays or indirectly in loops via top-charm associated production. In this letter, we shall investigate the latter case in the FCNC process

$$e^+e^- \rightarrow \bar{t}c \quad (\text{or } t\bar{c}). \quad (1)$$

Comparing t quark rare decays where the momentum transfer q^2 is limited, this process allows the large momentum transfer, which is actually determined by the energies available at LC. The process $e^+e^- \rightarrow \bar{t}c$ has some advantages for probing higher dimension operators at large momenta and striking kinetic signatures which are straight forward to detect in the clean environment of LC. In models beyond SM which can induce FCNC, there are

large underlying mass scales and large momentum transfer. So that these models are more naturally probed via top-charm ($t\bar{c}$) associated production than top quark rare decays[3].

The production cross section of the FCNC process $e^+e^- \rightarrow \bar{t}c$ in the SM has been calculated in Ref.[4] and a lot of theoretical activity involving top-charm production has been given within specific popular models beyond the SM. For example, studies of $\bar{t}c$ production in multi Higgs doublets models[5], in supersymmetry with R-Parity violation[6], in models with extra vector-like quarks[7], in technicolor models[8], and in a model independent approach[3]. The aim of this letter is to point out that the FC couplings predicted by topcolor-assisted technicolor(TC2) models[9] can give significant contributions to the FCNC process $e^+e^- \rightarrow \bar{t}c$, which may be detected in the future LC experiments.

An important issue in high-energy physics is to understand the mechanism of the mass generation. Given the large value of the top quark mass and the sizable splitting between the masses of the top and bottom quarks, it is natural to wonder whether m_t has a different origin from the masses of the other quarks and leptons and there may be a common origin for EWSB and top quark mass generation. Much theoretical work has been carried out in connection to the top quark and EWSB. The TC2 models[9], the top see-saw models[10] and the flavor universal TC2 models[11] are three of such examples. Such type of models generally predict a number of scalars with large Yukawa couplings to the third generation. For example, TC2 models predict the existence of scalars including the technipions ($\pi^0, \pi^\pm; \pi^{0,a}, \pi^{\pm,a}$) in the technicolor sector, and the top-pions ($\pi_t^0, \pi_t^\pm$) and the top-Higgs boson h_t^0 in the topcolor sector. Ref.[8] has considered the contributions of the scalars predicted by the technicolor model with a massless scalar doublet[12]. The results show that the cross section of the FCNC process $e^+e^- \rightarrow \bar{t}c$ is of the order of $10^{-3} - 10^{-5} fb$, which is too small to be detected in the future LC experiments. Further, Ref.[8] only considered the contributions of charged scalars to this process via the coupling $s^\pm u_i d_j$.

It is an important feature of TC2 models that the neutral scalars can induce the tree-level FC scalar couplings. It has been shown that the FC scalar couplings can give distinct new flavor-mixing phenomena which may be tested at both low- and high-energy

experiments[13, 14]. Thus, in this letter, we will mainly consider the contributions of the FC scalar coupling $s\bar{t}c$ (s is π_t^0 or h_t^0) to the FCNC process $e^+e^- \rightarrow \bar{t}c$ in the context of TC2 models and see whether the signals of TC2 models can be tested via the process $e^+e^- \rightarrow \bar{t}c$ in the future LC experiments. Our results show that the contributions of charged scalars and the topcolor gauge bosons to the FCNC process $e^+e^- \rightarrow \bar{t}c$ are all negligibly small. While the neutral scalars can give significant contributions which may be detected in the future LC experiments.

For TC2 models, the underlying interactions, topcolor interactions, are nonuniversal and therefore do not possess a GIM mechanism. This is an essential feature of this kind of models due to the need to single out the top quark for condensate. This nonuniversal gauge interactions result in FCNC vertices when one writes the interactions in the quark mass eigenbasis. Thus, the top-pions have large Yukawa coupling to the third family quarks and can induce the new FC scalar couplings including the $t - c$ transitions for the neutral top-pion π_t^0 [9, 13]:

$$\frac{m_t}{\sqrt{2}F_t} \frac{\sqrt{\nu_w^2 - F_t^2}}{\nu_w} \left[iK_{UR}^{tt}K_{UL}^{tt*}\bar{t}_L t_R \pi_t^0 + \sqrt{2}K_{UR}^{tt*}K_{DL}^{bb}\bar{t}_R b_L \pi_t^+ + \right. \\ \left. iK_{UR}^{tc}K_{UL}^{tt*}\bar{t}_L c_R \pi_t^0 + \sqrt{2}K_{UR}^{tc*}K_{DL}^{bb}\bar{c}_R b_L \pi_t^+ + h.c. \right], \quad (2)$$

where F_t is the top-pion decay constant and $\nu_w = \nu/\sqrt{2} = 174 GeV$. It has been shown that the values of coupling parameters can be taken as:

$$K_{UL}^{tt} = K_{DL}^{bb} = 1, \quad K_{UR}^{tt} = 1 - \epsilon, \quad K_{UR}^{tc} \leq \sqrt{2\epsilon - \epsilon^2}, \quad (3)$$

with a model-dependent parameter $\epsilon \ll 1$. In the following calculation, we will take $K_{UR}^{tc} = \sqrt{2\epsilon - \epsilon^2}$ and take ϵ as a free parameter.

The relevant Feynman diagrams for the contributions of the neutral top-pion π_t^0 to the process $e^+e^- \rightarrow \bar{t}c$ via the FC coupling are shown in Fig.1. Using Eq.(2) and other relevant Feynman rules, we can give the effective vertices $Z\bar{t}c$ and $\gamma\bar{t}c$ arising from π_t^0 :

$$\Lambda_{Z\bar{t}c}^\mu = ie \left[\gamma^\mu (F_{1Z} + F_{2Z}\gamma^5) + p_t^\mu (F_{3Z} + F_{4Z}\gamma^5) + p_c^\mu (F_{5Z} + F_{6Z}\gamma^5) \right], \quad (4)$$

$$\Lambda_{\gamma\bar{t}c}^\mu = \Lambda_{Z\bar{t}c}^\mu|_{F_{iZ} \rightarrow F_{i\gamma}}, \quad F_{i\gamma} = F_{iZ}|_{v_t=\frac{2}{3}, a_t=0}. \quad (5)$$

The form factors F_{iV} are expressed in terms of two- and three-point scalar integrals[16]. The expressions of F_{iV} are given in Appendix.

The total cross section can be written as:

$$\sigma = \frac{s - m_t^2}{32\pi s^2} \int_0^\pi \overline{|M|^2} \sin \theta d\theta, \quad (6)$$

with

$$\overline{|M|^2} = \frac{1}{4}|M|_Z^2 + \frac{1}{4}|M|_\gamma^2 + \frac{1}{2}ReM_Z M_\gamma^+. \quad (7)$$

The production cross section σ contributed by the π_t^0 is plotted in Fig.2 as a function of the top-pion mass m_{π_t} for the center-of-mass energy $\sqrt{s} = 500GeV$ and three of the parameter ϵ . From Fig.2 we can see that the FC scalar coupling $\pi_t^0 \bar{t}c$ could indeed give significant contributions to the process $e^+e^- \rightarrow \bar{t}c$. The value of the production cross section σ increase with ϵ and m_{π_t} increasing. For $m_{\pi_t} = 300GeV$, the cross section σ varies between $0.05fb$ and $0.12fb$ for the parameter ϵ in the range of $0.03 - 0.08$. For $\epsilon = 0.05$, σ varies between $0.076fb$ and $0.105fb$ for the top-pion mass m_{π_t} in the range of $200GeV - 450GeV$.

To see the effects of the center-of-mass energy on the production cross sections, we plot the σ versus $\sqrt{s}$ in Fig.3 for $m_{\pi_t} = 300GeV$ and three values of the parameter ϵ . One can see from Fig.3 that σ increases with $\sqrt{s}$ increasing for $\sqrt{s} \leq 480GeV$ and decreases with $\sqrt{s}$ increasing for $\sqrt{s} \geq 480GeV$. For $\sqrt{s} = 480GeV$, the production cross section σ reaches the maximum value $\sigma_{max} = 0.122fb$.

TC2 models also predict the existence of the neutral CP-even state, called top-Higgs boson h_t^0 , which is a $t\bar{t}$ bound and analogous to the σ particle in low energy QCD. Its mass can be estimated in the Nambu- Jona-Lasinio (NJL) model in the large N_C approximation and is found to be of the order of $m_{h_t} \approx 2m_t$ [13]. The main difference between π_t^0 and h_t^0 is that h_t^0 can couple to gauge boson pairs WW and ZZ at the tree-level, which is similar to that of the SM Higgs boson H^0 . However, the coupling coefficients of the couplings $h_t^0 WW$ and $h_t^0 ZZ$ are suppressed by the factor $\frac{F_t}{\nu_w}$ with respect to that of H^0 . Thus, the contributions of the top-Higgs boson h_t^0 to the FCNC process $e^+e^- \rightarrow \bar{t}c$ are similar to that of π_t^0 . Our numerical results are shown in Fig.4 for $m_{h_t} = 300GeV$.

From Fig.4 we can see that σ varies between $0.042fb$ and $0.184fb$ for $\sqrt{s}$ in the range of $300GeV - 1000GeV$, which is larger than the production cross section generated by π_t^0 . Combining the contributions of h_t^0 to the cross section σ with that of the top-pion π_t^0 , the total cross section σ_{total} given by the neutral scalars is in the range of $0.014fb - 0.35fb$ in most of the parameter space in TC2 models. For $\epsilon = 0.08$, $\sqrt{s} = 480GeV$ and $m_{\pi_t} = m_{h_t} = 450GeV$, the maximum value of σ_{total} can reach $0.35fb$.

The possible observable final states for the FCNC process $e^+e^- \rightarrow \bar{t}c$ are $\bar{b}cj_1j_2$, where j_1 and j_2 are light jets coming from $t \rightarrow bW^+$ followed by $W^+ \rightarrow u\bar{d}$ or $c\bar{s}$, and $b\bar{c}l^+\nu_l$, which comes from $t \rightarrow bW^+ \rightarrow bl^+\nu_l$ with $l = e, \mu$ or τ . These two final states occur with branching of $2/3$ and $1/3$, respectively. The leading SM background of the FCNC process $e^+e^- \rightarrow \bar{t}c$ mainly comes from W pair production as well as from W bremsstrahlung in $e^+e^- \rightarrow W + 2\text{-jets}$ [3]. We define the background-free observable cross section $\sigma_{\bar{t}c}^{\text{obs}}$ as the effective cross section including b -tagging efficiency (ϵ_b) and top quark reconstruction efficiency (ϵ_t): $\sigma_{\bar{t}c}^{\text{obs}} = \epsilon_b\epsilon_t\sigma_{\bar{t}c}$.

In order to estimate the number of the observable events, we assume $\epsilon_b = 60\%$ and $\epsilon_t = 80\%$ as done by Bar-Shalom and J. Wudka in Ref[3]. We consider two future e^+e^- collider scenarios: a LC with $\sqrt{s} = 500GeV$ and a yearly integrated luminosity of $L = 100fb^{-1}$ and a LC with $\sqrt{s} = 1000GeV$ and $L = 200fb^{-1}$. The yearly observable production events can be easily calculated. Our numerical results are shown in Fig.5. In Fig.5 we have assumed $m_{\pi_t} \approx m_{h_t} = m$ and take $\epsilon = 0.08$. We can see from Fig.5 that there will be several tens of the observable $\bar{t}c$ events to be generated in the future LC experiments. The number of the observable $\bar{t}c$ events generated in a LC with $\sqrt{s} = 500GeV$ is larger than that generated in a LC with $\sqrt{s} = 1000GeV$. Furthermore, for $350GeV < m < 500GeV$, the number generated in a LC with $\sqrt{s} = 500GeV$ increases with m increasing. Thus, it is likely that the signals of TC2 models can be more easily detected at a LC with $\sqrt{s} = 500GeV$ than at a LC with $\sqrt{s} = 1000GeV$.

If the masses of the neutral scalars are smaller than $350GeV$, $\bar{t}c$ and gg are the dominant decay modes of the scalars π_t^0 and h_t^0 . In this case, these new particles have significant contributions to the top-charm associated production at hadron colliders. This has been

extensively studied by Burdman[13]. His results show that the production cross section of the process $gg \rightarrow \pi_t^0(h_t^0) \rightarrow \bar{t}c$ is very large. There will be several thousands of $\bar{t}c$ events to be generated at the LHC, which are larger than the number of $\bar{t}c$ events produced at the future LC experiments. Thus, it is possible that the signals of TC2 models can be more easily detected at the LHC than at the LC experiments. However, the cross section of the process $gg \rightarrow \pi_t^0(h_t^0) \rightarrow \bar{t}c$ decrease by two or even more orders for $m_{h_t(\pi_t)} > 350\text{GeV}$. Furthermore, we must separate the signals from the large background before observation of the neutral scalars at the LHC. Thus, it is possible that the neutral scalars are more easily detected at the LC than at the LHC for $m_{h_t(\pi_t)} > 400\text{GeV}$.

To solve the phenomenological difficulties of the traditional TC models, TC2 models[9] were proposed by combining technicolor interactions with the topcolor interactions for the third generation at the scale about 1TeV . Thus, TC2 models predict number of technipions in the technicolor sector. These new particles also have contributions to the FCNC process $e^+e^- \rightarrow \bar{t}c$ via the flavor diagonal couplings $s^\pm u_i d_j$. The graphs of $e^+e^- \rightarrow \bar{t}c$ through charged technipions and top-pions loop are shown in Fig.6. Our calculation results show that the contributions of charged scalars to the FCNC process $e^+e^- \rightarrow \bar{t}c$ are smaller than that of the neutral scalars. In the most of parameter space, the production cross section is only in the range of $10^{-2} - 10^{-5}\text{fb}$, which can not be observed in the future LC experiments.

In TC2 models, to maintain electroweak symmetry between top and bottom quarks and yet to not generate $m_b \approx m_t$, the topcolor gauge group is usually taken to be a strongly coupled $SU(3) \times U(1)$. The $U(1)$ provides the difference that cause only top quark to condensate. Thus, TC2 models predict the existence of topcolor gauge bosons B_μ^A and an extra $U(1)_Y$ gauge boson Z' . Tree-level FCNC for the topcolor gauge bosons B_μ^A and Z' are generated when quarks fields are rotated to the mass eigenstate basis. The couplings of Z' are nonuniversal and stronger for the third generation, yielding potentially large top-charm couplings. Ref.[15] has shown that the FC coupling can indeed give significant contributions to the FCNC process $e^+e^- \rightarrow \bar{t}c$. For $M_{Z'} = M_{B'} = 1\text{TeV}$, the production cross section is of the order of $10^0 - 10^{-1}\text{fb}$. However, Ref.[17] has pointed out $B\bar{B}$

mixing provides strong lower bounds on the masses of B_μ^A and Z' bosons, i.e. there must be larger than 4 TeV . In this case, the cross section σ of the process $e^+e^- \rightarrow \bar{t}c$ given by topcolor gauge bosons is smaller than $10^{-2}fb$. Recently, E. H. Simmons[18] has shown that the B-meson mixing can also give lower bound on the mass of the Z' boson predicted by the flavor-universal TC2 models[11] or non-commuting extended technicolor(NCETC) models[19]. However, the bound is much weak, i.e. $M_{Z'} > 373GeV$. If we assume the mass of Z' predicted by the flavor-universal TC2 or NCETC models approximately equals to $500GeV$, then the cross section σ given by Z' is of the order of $10^0 - 10^{-1}fb$ in most of the parameter space. Thus, the Z' gauge boson predicted by these models can give significant contributions to the FCNC process $e^+e^- \rightarrow \bar{t}c$, which may be detected in the future LC experiments.

To completely avoid the problems, such as triviality and unnaturalness arising from the elementary Higgs field, various kinds of dynamical EWSB mechanisms have been proposed and among which TC2 theory is an attractive idea. TC2 models predict number of new scalars and gauge bosons. In this letter, we calculated the contributions of these new particles to the FCNC process $e^+e^- \rightarrow \bar{t}c$ in the framework of TC2 models. Our results shown that the contributions of charged scalars and topcolor gauge bosons to this process is very small. However, the neutral top-pion π_t^0 and top-Higgs h_t^0 can give significant contributions to the FCNC process $e^+e^- \rightarrow \bar{t}c$. With reasonable value of the parameters, the production cross section σ can reach $0.35fb$. There will be several tens of the observable $\bar{t}c$ events to be generated in the LC experiments with $\sqrt{s} = 500GeV$. Thus, the possible signals of π_t^0 and h_t^0 may be observed in the future LC experiments via this process.

Appendix: The form factors in the effective vertices $Z\bar{t}c$ and $\gamma\bar{t}c$ for the neutral top-pion π_t^0 :

$$F_{1Z} = g[B_0 + m_{\pi_t}^2 C_0 - 2C_{24} + m_t^2(C_{11} - C_{12}) - B_0^* - B_1']v_t,$$

$$F_{2Z} = g[B_0 + m_{\pi_t}^2 C_0 - 2C_{24} - m_t^2(C_{11} - C_{12}) + B_0^* + B_1' - 4\overline{C_{24}}]a_t,$$

$$F_{3Z} = 2m_t g(C_{21} + C_{22} - 2C_{23})v_t,$$

$$F_{4Z} = 2m_t g(-C_{21} - C_{22} + 2C_{23} - 2\overline{C_{22}} + \overline{C_{12}} + \overline{C_0})a_t,$$

$$F_{5Z} = 2m_t g(C_{22} - C_{23} + C_{12})v_t,$$

$$F_{6Z} = 2m_t g(-C_{22} + C_{23} + C_{12} - 2\overline{C_{22}} + 3\overline{C_{12}} + 2\overline{C_{23}} - 2\overline{C_{11}} - \overline{C_0})a_t.$$

Here $g = \frac{1}{16\pi^2} [\frac{m_t}{\sqrt{2}F_t} \frac{\sqrt{\nu_w^2 - F_t^2}}{\nu_w}]^2 K_{UR}^{tc} K_{UL}^{tt*}$, $v_t = \frac{1}{4s_W c_W} (1 - \frac{8}{3} S_W^2)$, $a_t = \frac{1}{4s_W c_W}$. The expressions of two- and three-point scalar integrals B_n and C_{ij} in this paper are[16]:

$$B_n = B_n(-\sqrt{s}, m_t, m_t), \quad B_n^* = B_n(-p_c, m_{\pi_t}, m_t),$$

$$B_n' = B_n(-p_t, m_{\pi_t}, m_t), \quad C_{ij} = C_{ij}(p_t, -\sqrt{s}, m_{\pi_t}, m_t, m_t),$$

$$C_0 = C_0(p_t, -\sqrt{s}, m_{\pi_t}, m_t, m_t), \quad \overline{C_{ij}} = C_{ij}(-p_c, \sqrt{s}, m_t, m_{\pi_t}, m_{\pi_t}),$$

$$\overline{C_0} = C_0(-p_c, \sqrt{s}, m_t, m_{\pi_t}, m_{\pi_t}).$$

Figure captions

Fig.1: Feynman diagrams for the contributions of the neutral top-pion π_t^0 to the process $e^+e^- \rightarrow \bar{t}c$. The dashed lines are the top-pion inner lines.

Fig.2: The production cross section σ contributed by π_t^0 as a function of the mass m_{π_t} for $\sqrt{s} = 500\text{GeV}$ and $\epsilon = 0.03$ (solid line), $\epsilon = 0.05$ (short dashed line), $\epsilon = 0.08$ (dot-dashed line).

Fig.3: The production cross section σ contributed by π_t^0 as a function of $\sqrt{s}$ for $m_{\pi_t} = 300\text{GeV}$ and $\epsilon = 0.03$ (solid line), $\epsilon = 0.05$ (short dashed line), $\epsilon = 0.08$ (dot-dashed line).

Fig.4: The production cross section σ contributed by h_t^0 as a function of $\sqrt{s}$ for $m_{h_t} = 300\text{GeV}$ and $\epsilon = 0.03$ (solid line), $\epsilon = 0.05$ (short dashed line), $\epsilon = 0.08$ (dot-dashed line).

Fig.5: The yearly observable total events contributed by h_t^0 and π_t^0 as a function of m for $\epsilon = 0.08$ and $\sqrt{s} = 500\text{GeV}$ (solid line), $\sqrt{s} = 1000\text{GeV}$ (short-dashed line). Here we assume $m_{h_t} = m_{\pi_t} = m$.

Fig.6: Feynman diagrams for the contributions of the charged technipions and top-pions to the process $e^+e^- \rightarrow \bar{t}c$. The dashed lines are the charged technipions or top-pions inner lines.

References

- [1] M. Beneke, et al. and A. Ahmadov, et al., “Top quark physics”, hep-ph/0003033; E. H. Simmons, “Top physics”, hep-ph/0011244.
- [2] For example, American Linear Collider Working Group, “Linear Collider Physics Resource Book for Snowmass 2001”, hep-ex/0106043.
- [3] Tao Han and J. L. Hewett, *Phys. Rev. D***60** (1999)074015; S. Bar-Shalom and J. Wudka, *Phys. Rev. D***60**(1999)094016.
- [4] For recent paper about the process $e^+e^- \rightarrow \bar{t}c$ in the standard model, see e.g., Chao-Shang Huang, Xiao-Hong Wu and Shou-Hua Zhu, *Phys. Lett. B***452**(1999)143.
- [5] W.-S. Hou and G.-L. Lin, *Phys. Lett. B***379**(1996)261; D. Atwood, L. Reina and A. Soni, *Phys. Rev. D***55**(1997)3156; S. Bar-Shalom, et al., *Phys. Rev. Lett.* **79**(1997)1217; *ibid.* *Phys. Rev. D***57**(1998)2957; J. Yi, et al, *Phys. Rev. D***57**(1998)4343.
- [6] U. Mahanta and H. Ghosal, *Phys. Rev. D***57**(1998) 1735; M. Chemtob and G. Moreau, *Phys. Rev. D***59**(1999)116012.
- [7] F. del Aguila, J. A. Aguilar-Saavedra and R. Miquel, *Phys. Rev. Lett.* **82**(1999)1628.
- [8] Xue-Lei Wang, et al., *Commun. Theor. Phys.* **27**(1997)325.
- [9] C. T. Hill, *Phys. Lett. B***345**(1995)483; K. Lane, and T. Eichten, *Phys. Lett. B***352**(1995)382; K. Lane, *Phys. Lett. B***433**(1998)96; G. Cvetič, *Rev. Mod. Phys.* **71**(1999)513.
- [10] B. A. Dobrescu and C. T. Hill, *Phys. Rev. Lett.* **81**(1998)2634; R. S. Chivukula, B. A. Dobrescu, H. Georgi and C. T. Hill, *Phys. Rev. D***59**(1999)075003.
- [11] M. B. Popovic and E. H. Simmons, *Phys. Rev. D***58**(1998)095007.
- [12] C. D. Carone and H. Georgi, *Phys. Rev. D***49**(1994)1427.

- [13] H.-J. He and C.-P. Yuan, *Phys. Rev. Lett.* **83**(1999)28; G. Burdman, *Phys. Rev. Lett.* **83**(1999)2888.
- [14] Chongxing Yue, et al., *Phys. Lett. B***509**(2001)294; Chongxing Yue, et al., *Phys. Rev. D***63**(2001)115002; Chongxing Yue, et al., *Phys. Rev. D***64**(2001)095004.
- [15] Chongxing Yue and Weibin Li, *Mod. Phys. Lett. A***15**(2000)361.
- [16] G. Passarino and M. Veltman, *Nucl. Phys. B***160**(1979)151; A. Axelrod, *ibid.* **B209**(1982)349; M.Clements, et al., *Phys. Rev. D***27**(1983)570.
- [17] G. Burdman, K. Lane and T. Rador, *Phys. Lett. B* **514**(2001)41.
- [18] E. H. Simmons, “Limitations of B-meson mixing bounds on technicolor theories”, hep-ph/0111032.
- [19] R. S. Chivukula, E. H. Simmons and J. Terning, *Phys. Lett. B***331**(1994)383.




Fig.1




Fig.2




Fig.3




Fig.4




Fig.5




Fig.6